

適応疾患/悪性リンパ腫

プロトコール名 **P-L-1 / R-CHOP 療法**

| 使用薬剤 | 備考 | 投与方法 | 投与日 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 1 | 2 | 3 | 4 | 5 | 6 | |
|--|--|------|-----|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|---|---|---|---|---|---|---|
| | | | Day | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 抗ヒスタミン薬 | day1 | 経口 | | ↓ | | | | | | | | | | | | | | | | | | | | | ↓ | | | | | | |
| 解熱鎮痛剤 | day1 | 経口 | | ↓ | | | | | | | | | | | | | | | | | | | | | ↓ | | | | | | |
| ① ソルコーテフ200mg 生食100ml | 30分 | 点滴静注 | | ↓ | | | | | | | | | | | | | | | | | | | | | ↓ | | | | | | |
| ② リツキサ 375mg/m ² /day 生食(適宜) ml | 25ml/h で 1h 100ml/h で 1h 以後200ml/h | 点滴静注 | | ↓ | | | | | | | | | | | | | | | | | | | | | ↓ | | | | | | |
| | | | Day | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| プレドニン 100mg/body/day | day3-7 | 経口 | | | | ↓ | ↓ | ↓ | ↓ | ↓ | | | | | | | | | | | | | | | | | | ↓ | ↓ | ↓ | ↓ |
| ① グラニセトロンバッグ | 30分 | 点滴静注 | | | | ↓ | | | | | | | | | | | | | | | | | | | | | ↓ | | | | |
| ② アドリアシン 50mg/m ² /day 生食100ml | 30分 | 点滴静注 | | | | ↓ | | | | | | | | | | | | | | | | | | | | | ↓ | | | | |
| ③ オンコビン 1.4mg/m ² /day 生食100ml | 最大2mg/body boius(全開で) | 点滴静注 | | | | ↓ | | | | | | | | | | | | | | | | | | | | | ↓ | | | | |
| ④ エンドキサ 750mg/m ² /day 生食500ml | 120分 | 点滴静注 | | | | ↓ | | | | | | | | | | | | | | | | | | | | | ↓ | | | | |
| ⑤ ビカーボン500ml | 適宜 | 点滴静注 | | | | ↓ | | | | | | | | | | | | | | | | | | | | | ↓ | | | | |

* 1コース28日

7

7

↓